

## Engineering Conferences International ECI Digital Archives

---

Cell Culture Engineering XV

Proceedings

---

Spring 5-12-2016

# A community genome-scale model of Chinese Hamster ovary cell metabolism identifies differences in the efficiency of resource utilization for various bioprocesses

Hooman Hefzi

*University of California - San Diego, [hhefzi@ucsd.edu](mailto:hhefzi@ucsd.edu)*

Follow this and additional works at: [http://dc.engconfintl.org/cellculture\\_xv](http://dc.engconfintl.org/cellculture_xv)



Part of the [Biomedical Engineering and Bioengineering Commons](#)

---

### Recommended Citation

Hooman Hefzi, "A community genome-scale model of Chinese Hamster ovary cell metabolism identifies differences in the efficiency of resource utilization for various bioprocesses" in "Cell Culture Engineering XV", Robert Kiss, Genentech Sarah Harcum, Clemson University Jeff Chalmers, Ohio State University Eds, ECI Symposium Series, (2016). [http://dc.engconfintl.org/cellculture\\_xv/187](http://dc.engconfintl.org/cellculture_xv/187)

This Abstract is brought to you for free and open access by the Proceedings at ECI Digital Archives. It has been accepted for inclusion in Cell Culture Engineering XV by an authorized administrator of ECI Digital Archives. For more information, please contact [franco@bepress.com](mailto:franco@bepress.com).

# **A COMMUNITY GENOME-SCALE MODEL OF CHINESE HAMSTER OVARY CELL METABOLISM IDENTIFIES DIFFERENCES IN THE EFFICIENCY OF RESOURCE UTILIZATION FOR VARIOUS BIOPROCESSES**

Hooman Hefzi\*, Department of Bioengineering, University of California, San Diego; Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark  
hhefzi@ucsd.edu

Kok Siong Ang\*, Department of Chemical and Biomolecular Engineering, National University of Singapore  
Michael Hanscho\*, Austrian Centre of Industrial Biotechnology

Michael Betenbaugh, Department of Chemical and Biomolecular Engineering, Johns Hopkins University

Mikael Andersen, Department of Systems Biology, Technical University of Denmark

Bernhard Palsson, Department of Bioengineering, University of California, San Diego; Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark

Lars Nielsen, Australian Institute for Bioengineering and Nanotechnology, The University of Queensland

Nicole Borth, Austrian Centre of Industrial Biotechnology; Institute for Applied Microbiology, Department of Biotechnology, University of Natural Resources and Life Sciences

Dong-Yup Lee, Department of Chemical and Biomolecular Engineering, National University of Singapore

Nathan Lewis, Department of Pediatrics, University of California, San Diego; Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark

\*These authors contributed equally to this work

**Key Words:** CHO, systems biology, modeling, metabolic engineering

Genome-scale models of metabolism have successfully been employed in many microbial and eukaryotic metabolic engineering efforts by guiding pathway engineering and media optimization. They have also been used to explore the genotype-phenotype relationship in mammalian cells. The publication of the genomic sequence for Chinese hamster ovary (CHO) cells has allowed generation of genome-scale metabolic models (GeMs) for this organism. Here we have developed a high-quality community CHO GeM via careful reconciliation and manual curation of three independently developed CHO GeMs. This metabolic model, consisting of over 4000 metabolites and 6000 reactions, is capable of integrating proteomic, transcriptomic, and metabolomic data and can accurately simulate experimentally measured growth rates. Integration of transcriptomic and proteomic data from CHO-K1 and CHO-S shed light on the enzymatic basis for various amino acid auxotrophies characteristic of the cell lines. We show that experimental arginine and cysteine auxotrophies are recapitulated by model predictions (via reaction inactivation) while the characteristic proline auxotrophy is not, due to detectable levels of expression in biosynthetic pathways for this amino acid. We additionally used the model to assess the metabolic limitations on recombinant protein producing lines subject to different cell line and process modifications and found that some alterations result in specific productivities up to 20-fold lower than computational predictions of metabolically feasible production rates. The results indicate a possible secretory bottleneck and implicate engineering the secretory pathway as a lucrative target to pursue in future CHO cell line engineering.